

## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <a href="http://about.jstor.org/participate-jstor/individuals/early-journal-content">http://about.jstor.org/participate-jstor/individuals/early-journal-content</a>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

armored dinosaurs from the Hell Creek Beds of Montana. material, however, upon which he bases his restoration was scanty—too scanty to serve as a satisfactory basis for a restoration—consisting of the skull, a number of vertebræ, a somewhat problematical scapula, and a number of dermal scutes. dangers of such attempted restorations as the present one on such imperfect material are apparent here, since Ankylosaurus is either very closely allied to or identical with Stegopelta Williston, a genus overlooked by Mr. Brown. And Stegopelta, which is represented by considerable material in the University of Chicago collections, is as closely allied as well may be with Polacanthus Hulke of England, and must go in the same family. A comparison of Nopsca's restoration of Polacanthus, which is essentially correct, save for the skull, will show the real form of Ankylosaurus, very different from that given by Mr. Brown. Stegopelta, moreover, has spines, and the body is covered in large part by small scutes, though there are also large ones, as figured by Mr. Brown; and the tail rings are correctly placed The writer has examined the specimen in the British Museum, and is assured of the relationships. Stegopelta will shortly be more fully described and figured by Dr. Moodie.

S. W. WILLISTON.

## PARASITOLOGY

The Evolution of Parasitism.—In a recent paper entitled: "The Influence of Symbiosis upon the Pathogenicity of Microorganisms (The Evolution of Parasitism)," Musgrave<sup>1</sup> has brought together some most important items in an extremely suggestive He defines symbiosis as representing all phases of association between living organisms, beginning with commensalism, on the one hand, and including true parasitism, on the other, in which either component is influenced in nutrition, metabolism, production or in some other manner by the presence of the other. While this use of the term is distinctly new and rather at variance with the older usage of van Beneden the meaning of the author is clear. He endeavors to show that symbiotic combinations between microorganisms are responsible for uninterpreted phenomena in the etiology and pathology of disease. more, he adduces evidence to show that changes is symbiosis may produce changes in metabolism and also, as a result of this,

<sup>&</sup>lt;sup>1</sup> Philippine Jour. Sc., B 3: 78.

changes in the pathogenic character of parasites and in the susceptibility of hosts. The work of other authors is cited indicating the influence that surroundings have on cultures of bacteria even under experimental conditions and leading to the conclusion that the relation between nutrition, metabolism and production is far more significant than has heretofore been considered in laboratory technique. If such influences modify artificial cultures, there will surely be a much greater effect from the complex conditions surrounding mixed cultures in unknown symbiosis.

Among animal parasites nutrition, metabolism and production certainly are more complicated than with bacteria, and unless this fact is constantly in mind the whole subject of parasitism is apt to be considered along too narrow lines and our specificities to be regarded as too exacting. The question, just as with bacteria, not only involves a study of the interaction between an animal host and animal parasites, but a study of host under constantly varying conditions being acted upon by parasites influenced by an ever-changing environment. A battle, as it were, takes place between two sets of very complex influences, leading to death or disease of the host on the one hand, to destruction or commensalism on the part of the parasite on the other, or finally quite frequently to what Theobald Smith has called a condition of balanced parasitism.

Most suggestive in this direction is the experimental work which has been done with amœba. Chief among the investigators are Musgrave and Clegg, who found that the cultivation of amœba could not be accomplished satisfactorily except in the presence of other living organisms, and this symbiosis is more or less specific although its specific character may be changed both experimentally and probably also under natural conditions.

The author sums up his paper as follows:

"I am convinced that such an evolution of parasitism from an amedic and bacterial symbiosis, in water or elsewhere in nature, through a beginning or mixed tissue and bacterial parasitism in amedic ulcers of the colon, to a true tissue parasitism in the internal organs of the body, or even to a blood invasion itself, is a matter of frequent occurrence. The most promising field for laboratory research in the future will be the study of cause and effect, in the complex relations in which they occur in nature, of the interrelation and interaction of microorganisms with each other and in their environment of complex symbiosis and the ever-changing and multiple conditions found in hosts."